arXiv:1704.03083v1 [math.GR] 10 Apr 2017

A characterization of ZM-groups

Marius Tărnăuceanu

April 11, 2017

Abstract

In this short note we give a characterization of ZM-groups that uses the functions defined and studied in [3, 4]. This leads to a proof of Conjecture 6 in [4].

MSC (2010): Primary 20D60, 11A25; Secondary 20D99, 11A99.
Key words: Gauss formula, Euler's totient function, finite group, order of an element, exponent of a group.

1 Introduction

Given a finite group G, we consider the functions

$$\varphi(G) = |\{a \in G \mid o(a) = \exp(G)\}| \text{ and } S(G) = \sum_{H \leq G} \varphi(H).$$

The class of finite groups G for which $S(G) = |G|$ has been partially determined in [4]. We are now able to complete this study by proving the following result.

Theorem 1.1. *Let G be a finite group. Then $S(G) \geq |G|$, and we have equality if and only if G is a ZM-group, i.e. a group with all Sylow subgroups cyclic.*

In particular, since a ZM-group is nilpotent if and only if it is cyclic, we obtain the result in Conjecture 6 of [4].

Corollary 1.2. *Let G be a finite nilpotent group. Then $S(G) \geq |G|$, and we have equality if and only if G is cyclic.*

2 Proof of Theorem 1.1

Let $C(G)$ be the poset of cyclic subgroups of G. For every divisor d of $|G|$ we denote by n_d the number of cyclic subgroups of order d of G and by n'_d the number of elements of order d in G. Then we have

$$n'_d = n_d\varphi(d)$$

because a cyclic subgroup of order d contains $\varphi(d)$ elements of order d. One obtains

$$S(G) \geq \sum_{H\in C(G)} \varphi(H) = \sum_{H\in C(G)} \varphi(|H|) = \sum_{d\,|\,|G|} n_d\varphi(d) = \sum_{d\,|\,|G|} n'_d = |G|,$$

as desired.

Assume now that $S(G) = |G|$. Then $\varphi(H) = 0$ for all non-cyclic subgroups H of G. Since for any p-group P we have $\varphi(P) \neq 0$ (see [3]), it follows that all Sylow subgroups of G are cyclic, that is G is a ZM-group.

Conversely, assume that G is a ZM-group. By [2] such a group is of type

$$\mathrm{ZM}(m,n,r) = \langle a, b \mid a^m = b^n = 1,\ b^{-1}ab = a^r\rangle,$$

where the triple (m,n,r) satisfies the conditions

$$\gcd(m,n) = \gcd(m,r-1) = 1 \text{ and } r^n \equiv 1 \pmod m.$$

It is clear that $|\mathrm{ZM}(m,n,r)| = mn$. The subgroups of $\mathrm{ZM}(m,n,r)$ have been completely described in [1]. Set

$$L = \left\{(m_1,n_1,s) \in \mathbb{N}^3 \ \mid\ m_1|m,\ n_1|n,\ s<m_1,\ m_1|s\frac{r^n-1}{r^{n_1}-1}\right\}.$$

Then there is a bijection between L and the subgroup lattice $L(\mathrm{ZM}(m,n,r))$ of $\mathrm{ZM}(m,n,r)$, namely the function that maps a triple $(m_1,n_1,s) \in L$ into the subgroup $H_{(m_1,n_1,s)}$ defined by

$$H_{(m_1,n_1,s)} = \bigcup_{k=1}^{\frac{n}{n_1}} \alpha(n_1,s)^k\langle a^{m_1}\rangle = \langle a^{m_1}, \alpha(n_1,s)\rangle,$$

where $\alpha(x,y) = b^xa^y$, for all $0 \leq x < n$ and $0 \leq y < m$. We can easily check that

$$|H_{(m_1,n_1,s)}| = \exp(H_{(m_1,n_1,s)}) = \frac{mn}{m_1n_1},$$

and so

$$\varphi(H_{(m_1,n_1,s)}) \neq 0 \text{ if and only if } H_{(m_1,n_1,s)} \text{ is cyclic.}$$

This shows that

$$S(G) = \sum_{H \in C(G)} \varphi(H) = |G|,$$

completing the proof. ∎

References

[1] W.C. Calhoun, *Counting subgroups of some finite groups*, Amer. Math. Monthly **94** (1987), 54-59.

[2] B. Huppert, *Endliche Gruppen*, I, Springer Verlag, Berlin, 1967.

[3] M. Tărnăuceanu, *A generalization of the Euler's totient function*, Asian-Eur. J. Math. **8** (2015), no. 4, article ID 1550087.

[4] M. Tărnăuceanu, *On a generalization of the Gauss formula*, Asian-Eur. J. Math. **10** (2017), no. 1, article ID 1750008.

Marius Tărnăuceanu
Faculty of Mathematics
"Al.I. Cuza" University
Iaşi, Romania
e-mail: tarnauc@uaic.ro